




NASO-JEJUNAL (NJ) TUBE

THE FOLLOWING INFORMATION IS AIMED AT SUPPORTING YOU ON HOW TO CARE FOR THE NJ TUBE.

IMPORTANT NOTE: It is important that you contact your Nutricia Homeward Nurse if you need additional training on the Flocare pump (if applicable) or tube feeding.



DAILY CARE OF THE NASO-JEJUNAL TUBE

It is important to keep the tube in good condition to avoid unnecessary replacement of the tube. Regular care and flushing of the tube will help to prevent the tube becoming blocked.

- Wash hands before and after handling the tube
- Confirm the position of the tube by checking the cm graduation marking at the nostril remains the same cm marking made after initial placement in hospital. Position must be checked before administration of any water, enteral nutrition or medication
- Flush the tube as per training provided by Nutricia Homeward Nurse, using a 60ml enteral syringe with water (type and volume as recommended for jejunal use by the managing healthcare professional) to prevent the tube from blocking. Water should be administered before and after enteral nutrition/medication administration or every 4-6 hours if feeding is not in progress (except during the night) or as recommended by the managing healthcare professional
- Care of the nose: fixation tape should be changed at least weekly, but more frequently if it loosens or becomes dirty. When changing the tape, clean skin thoroughly and try to alter the position of the tape from where it was last fixed. If skin or the nostril becomes sore or red contact the managing healthcare professional for instructions on how to proceed. Avoid the use of cream or powders as they can damage the tube
- Never attempt to replace the tube as this type of tube can only be replaced by the healthcare professional in the hospital
- Check the tube daily to ensure it is intact and that the tube is not kinked or blocked and remains securely fixed to the face.

HOW DO I CHECK THE POSITION OF THE NASO-JEJUNAL TUBE?

It is important to check the tube position prior to administering anything via the feeding tube, after a coughing fit or following a vomiting episode.

- Confirm the position of the tube by checking the cm graduation marking at the nostril remains the same cm marking made after initial placement
- Position must be checked before administration of any water, enteral nutrition or medication
- If there is any doubt about the position of the tube, do **not** use the tube and contact the managing healthcare professional for instructions on how to proceed.

IMPORTANT NOTES

It is important that the patient is sitting at a 45° angle during enteral feeding and for at least one hour after enteral nutrition has been administered.

You are advised to contact the managing healthcare professional immediately if there is any reason that medication or enteral nutrition needs to be delayed or omitted due to no available enteral or oral route.

TROUBLESHOOTING NASO-JEJUNAL TUBES

THE NASO-JEJUNAL TUBE HAS BECOME BLOCKED

If there is resistance when flushing the naso-jejunal tube, do not force water into the tube:

- Confirm tube remains in correct position before administering anything down the tube
- Using a gentle pull and push technique, flush the tube using a 60ml enteral syringe with warm water (type and volume as recommended for jejunal use by the managing healthcare professional). **Do not use acidic solutions such as fruit juices or cola as they can curdle the enteral nutrition product**
- If unable to clear the blockage contact the managing healthcare professional or Nutricia Homeward Nurse for further instructions.

THE NASO-JEJUNAL TUBE HAS COME OUT

It is important that a new naso-jejunal tube is placed in time for the next delivery of enteral nutrition or medication administration.

A naso-jejunal tube must always be replaced in a hospital setting therefore do not attempt to replace a new tube at home.

Follow the steps below:

- Remain calm
- Contact the managing healthcare professional and explain that the naso-jejunal tube has come out
- If the managing healthcare professional is unavailable contact the GP or hospital emergency department to let them know the tube will need to be replaced. This will give the department time to find the medical notes and to ensure a member of staff is available to replace the naso-jejunal tube
- Take the tube that has fallen out with you in a clean plastic bag so that the staff can identify which type of tube is required.

IMPORTANT: Always seek urgent medical advice if there are any signs of abdominal pain, chest pain or breathing difficulties.

The Nutricia Homeward Nursing Service advice literature is provided as guidance following Nutricia Homeward Nursing Service training, and should not be used as a substitute for medical advice. Always contact your healthcare professional prior to making any changes to your tube feeding regimen.

Please contact the Nutricia Homeward Nurse or managing healthcare professional for the most up to date version every year. Provided on behalf of Nutricia Ltd at the request of the NHS managing healthcare professional.

IMPORTANT: Always seek urgent medical advice if there are any signs of abdominal pain, chest pain or breathing difficulties.

Nutricia Ltd
White Horse Business Park, Trowbridge
Wiltshire BA14 0XQ
nutriciafocare.com
SCC3641-03/19



HOMEWARD
NURSING SERVICE